

Key

AP Chemistry Unit 6 Bonding Practice test Review

I. Directions: Answer each of the following questions. No calculators are allowed unless specified.

Goal: I can differentiate between an ionic bond, covalent bond or metallic bond.

For # 1-8, identify which statements below is a characteristic of an ionic bond(I), covalent bond(C) or a metallic bond(M).


- | | |
|--|---|
| <u>I</u> 1. Potassium iodide. | <u>M</u> 5. Conducts electricity in the solid state |
| <u>M</u> 2. Aluminum. | <u>C</u> 6. More sharing of electrons. |
| <u>I</u> 3. More transfer of electrons | <u>M</u> 7. Have mobile electrons. |
| <u>C</u> 4. Water | <u>C</u> 8. Carbon disulfide |

9. Which one of the following element combinations is likely to produce ionic bonds in a compound?

- a) Li and F metal & Nonmetal
 b) B and O 2 Nonmetals
 c) N and O 2 Nonmetals
 d) P and S 2 Nonmetals
 e) Cl and Br 2 Nonmetals

Goal: I can draw the Lewis symbols for any element on the periodic table.

10. Below each element, draw its Lewis dot structures




11. Which one of the following has eight valence electrons?

- a) Ti^{4+} *lose 4 ve, pull next level*
 b) Kr *lose 1 ve*
 c) Cl^-
 d) Na^+
 e) All of these!

Goal: I can define Lattice Energy and its relationship to ionic compounds and their melting points. I will also be able to describe the factors which affect lattice energy.

12. Explain, using the factors that affect lattice energy why the following compounds increase in lattice energy: $\text{KBr} < \text{KF} < \text{MgSe} < \text{MgO} < \text{AlN}$



13. Which of the following compounds would you expect to have the lowest melting point? **Smaller LE**

- a) NaF b) Na₂O c) NaBr d) NaI e) CaO

14. Calcium oxide has a lower melting point than magnesium oxide due to the

- a. greater charge on the Mg²⁺ than of Ca²⁺. **NO SAME!**
 b. greater charge on the Ca²⁺ than of Mg²⁺ " "
 c. the greater distance between the charges in CaO.
 d. the greater distance between the charges in MgO.
 e. equal charges on the Mg²⁺ than of Ca²⁺.

Ca has Larger Radius than Mg

15. In ionic bond formation, the lattice energy of ions _____ as the magnitude of the ion charges _____ and the radii _____.

- a. increases, decrease, increase.
 b. increases, increase, increase.
 c. decreases, decrease, increase.
 d. increases, increase, decrease.
 e. decreases, increase, decrease.

Goal: I can differentiate between a single, double, and a triple covalent bond in terms of number of shared electrons, length and strength.

16. Which of the following elements is **most** likely to participate in the formation of multiple bonds?

- a) H · b) Na · c) :Cl: d) S e) F:

17. Which molecule has the weakest bond?

- a) CO b) O₂ c) Cl₂ d) NO e) N₂
- C≡O O=O Cl-Cl N≡N

weakest = longest Single Bond

18. As the number of covalent bonds between two atoms increases, the distance between the atoms _____ and the strength of the bond between them _____.

- a. increases, increases.
 b. decreases, increases.
 c. increases, decreases.
 d. decreases, decreases.
 e. is unpredictable

Goal: I am able to define electronegativity and apply the electronegative trends to the periodic table.

19. In which of the following are the elements listed in order of increasing electronegativity?

- a. Ba, Zn, C, Cl
- b. O, N, S, Cl**
- c. N, P, As, Sb
- d. K, Ba, Si, Ga
- e. Li, K, Na, Ca

Goal: I am able to predict the relative polarities of bonds using either the periodic table or electronegativity values. I can also describe the factors which affect bond polarity and dipole moment.

20. Which of the following bonds is expected to be the most polar?







- a. C-Si $2.5 - 1.8 = .7$
- b. C-N $2.5 - 3 = .5$
- c. O-C $3.5 - 2.5 = 1.0$**
- d. S-C $2.5 - 2.5 = 0$
- e. H-C $2.1 - 2.5 = .4$

21. C_2O C_2F N_2O K_2Br S_2O Rb_2Cl
 The bond above that is the most polar is N_2O and the least polar is Rb_2Cl .

- a. C, O K, Br
- b. Rb, Cl F, N, O**
- c. C, F S, O
- d. Rb, Cl S, O
- e. N, O C, F

Goal: Write the Lewis structures for molecules and ions containing covalent bonds using the periodic table.



22. Draw Lewis dot structures for the following molecules:



Goal 8: Explain the concept of resonance and draw resonance structures for molecules or polyatomic ions.


23. Which molecule below possesses resonance?

- a) BCl_3 b) SO_2 c) CO_2 d) BeF_2 e) SO_4^{2-}



Questions 24-26: Consider the chemical bonds found in the white solid, potassium hydrogen phosphate, K_2HPO_4 . For each bond specified, choose the best description from the list of bond types below.

- a. ionic bond
- b. hydrogen bond
- c. single covalent bond
- d. double covalent bond
- e. covalent bond with resonance



24. phosphorus/oxygen bond C P-O

25. potassium/hydrogen phosphate bond A $\text{K}^+ \text{HPO}_4^{2-}$


26. oxygen/hydrogen bond C O-H

Goal 9: Use the concept of formal charge to predict the most stable resonance structure.

27. The formal charge on the nitrogen in a nitrite ion (NO_2^-) is

- a) -2 b) -1 c) 0 d) 1 e) 2

$N = 1 \times 5 = 5$
 $O = 2 \times 6 = 12$
 $2e = 14$




28. Identify all the following statements that are TRUE about formal charge.

T a. The formal charge on an atom is determined by taking the valence electrons minus the electrons assigned to that atom.

F b. A Lewis structure is valid if a negative formal charge is on the least electronegative element.

F c. The formal charge on the oxygen in a water molecule is -2.



F d. For neutral compounds, the formal charges add up to various numbers and for ions the formal charges add up to equal the charge on the ion.

F e. Lewis structures with large formal charges are most likely to exist.

small


Goal : Describe the 3 common exceptions to the octet rule and provide examples of each.

The three common exceptions are:

- A. Molecules that have an odd number of electrons.
- B. Molecules in which there is less than an octet of electrons.
- C. Molecules in which there is more than an octet of electrons.

Knowing this, identify the exception that each molecule best fits.

29. XeF₄




30. NO


$$\begin{array}{r} N \ 1 \times 5 = 5 \\ O \ 1 \times 6 = 6 \\ \hline 11 \end{array}$$



31. BH₂Cl




32. SeF₄



33. ClO₂⁻

$$\begin{array}{r} Cl \ 1 \times 7 = 7 \\ O \ 2 \times 6 = 12 \\ + 1 \times 1 \\ \hline 20 \end{array}$$



34. Which of the following is most likely to form compounds involving an expanded octet?


- a) Li
No
d
- b) N
No
d sub shell
- c) F
No
d
- d) Ne
No
d
- e) S
has d sub shell

Target 11: Relate bond enthalpies to bond strengths and use bond enthalpies to estimate ΔH for reactions.

35. Bond enthalpy is _____ positive and is defined as the energy _____ to break a bond.


- a) always, required
- b. never, required
- c. sometimes, produced
- d. always, produced
- e. never, required

36. Given the bond dissociation energies below, calculate the standard molar enthalpy of formation of ClF_3 .



Bond	Dissociation Energy (kJ/mol)
Cl-Cl	243
F-F	159
Cl-F	255

Cl 117 = 7
F 817 = 21
298 e



- a) 210 kJ/mol b) 147 kJ/mol c) -33 kJ/mol d) -405 kJ/mol **e) -810 kJ/mol**

$$\begin{aligned} \Delta H &= \sum \text{Bonds Broken} - \sum \text{Bonds Formed} \\ &= (\text{Cl-Cl}) + 3(\text{F-F}) - [2(3 \times \text{Cl-F})] \\ &= 243 + 3(159) - (6(255)) \text{ kJ/mol} \\ &= 720 \frac{\text{kJ}}{\text{mol}} - 1530 \frac{\text{kJ}}{\text{mol}} \\ &= -810 \frac{\text{kJ}}{\text{mol}} \end{aligned}$$

Goal: I can determine the bond angle, geometric shape and the electron domain shape of a molecule or ion based upon the V.S.E.P.R. Theory.

37. Which of the following has a nonbonding pair of electrons on the central atom?



- a. BCl_3 **b. NH_3** c. CCl_2Br_2 d. PF_5 e. SO_4^{2-}

~ 1 lone pair




38. Which one of the following may we draw both polar and nonpolar Lewis structures?

- a. CHCl_3
b. NH_3
c. BF_3
d. SF_2Cl_4
e. SO_2






39. Which one of the following is NOT a linear structure?

- a. I_2 b. I_2^- c. CO_2 **d. H_2S** e. $\text{H}-\text{C}\equiv\text{C}-\text{H}$



40. The Lewis structure of the cyanide ion most closely resembles which of the following?

- a. N_2** b. O_2 c. CO_2 d. NO e. C_2H_2




41. Which angle is NOT expected in any molecule?

- a. 60°
- b. 90°
- c. 109.5°
- d. 120°
- e. All of these are reasonable angles.

Goal: I can explain why nonbonding electron domains exert a greater repulsive interaction on other domains than do bonding electron domains.

42. Which molecule below would have the greatest repulsion of electrons resulting in the smallest bond angle.




Goal: I can predict from the shape of a molecule whether it is polar or nonpolar.

43. Which of the bonds below is the most polar? (Review from Chapter 8)


Element	Si	H	C	S	N	O
electronegativity	1.8	2.1	2.5	2.5	3.0	3.5

- a. C - Si b. C - N c. O - C d. S - C e. H - C
- Handwritten electronegativity differences: a. 2.5-1.8=0.7; b. 2.5-2.1=0.4; c. 2.5-3.5=1.0; d. 2.5-2.5=0; e. 2.1-2.5=0.4*

44. How many of the molecules below are polar. (Circle answers that are polar!)




45. Which one molecule below will have a zero dipole moment?



Goal: I can explain the concept of hybridization and its relationship to geometrical structure. I can predict the type of hybrid orbitals of an atom in a molecule.

46. The SF₅⁻ ion has a square pyramid structure. The hybridization of the orbitals in sulfur is:





47. Sulfur forms the following compounds: SO_2 , SF_6 , SCl_4 , SCl_2 . Which form of hybridization is NOT represented by these molecules?

- a. sp b. sp^2 c. sp^3 d. dsp^3 e. d^2sp^3

↳ Sulfur will always need more than 2 bonds




Goal: I can explain the difference between a sigma bond and a pi bond. I can also determine the number of sigma bonds and pi bonds in a molecule or ion.

48. For # a - f, identify which statements below is a characteristic of a pi (p) or a sigma (s) bond.

- S a. Formed by the head-to-head bond between atoms. 
- P b. Formed above and below the bond axis. 
- P c. A triple bond would contain two of these.
- P d. This bonding occurs from the sideways overlap of an electron in p orbitals.
- S e. A triple bond would contain ~~two~~ one of these.

49. Which of the following has the fewest pi bonds and is nonpolar?

- a. HCCH
b. CO_2
c. CO_3^{2-}
d. N_2
e. SO_2




50. Which one of the following is true when the $\text{C}=\text{C}$ and $\text{C} \equiv \text{C}$ bonds are compared?

- T (a) The triple bond is shorter than the double bond.
b. The double bond contains more pi bonds. No Triple has 2, = has 1
c. The double-bond energy is higher than the triple-bond energy. No opposite
d. The double bond contains less sigma bonds.

No they both contain 1 sigma each

I. FREE RESPONSE: Complete each of the following free response questions in the spaces provided.

1. Draw the Lewis dot structure for a nonpolar molecule that has a molecular formula of $\text{Si}_2\text{H}_2\text{Br}_2$. (Si is double bond to Si)




a. There are 5 sigma bonds and 1 pi bond(s) in this molecule?


b. The hybridization around each ^{silicon}carbon is SP^2 , and the bond angle is 120° .

This molecule is nonpolar because it is Sym and it has _____ dipoles.

c. Draw the two polar isomers of this compound and indicate where the dipole is located.




2. Draw the structure for acetic acid ($\text{H}_3\text{C}-\text{COOH}$) below. NOTE: The second carbon is double bonded to the first oxygen and single-bonded to the second O. The H is attached to the second oxygen.



a. What is the molecular geometry around each carbon atom?

Tetrahedral → trigonal planar

b. What type of hybrid orbitals does each carbon have?



3. Name three elements which . . .

a) cannot have expanded octets when it is the central atom in a molecule.

b) do not need an octet to form stable compounds.

Lewis

Molecule or Ion	Structure	e-domain geometry	Molecular geometry or shape	Bonded / Nonbonded e- domains	Polar or Nonpolar	Hybridizaion	Bond Angle
PH ₃ P 5 H 1x3=3 8ve		tetra	Trigonal planar	3 Bonds 1 Lone pair	Polar	3 sigma 1 lone pair sp ³	109°
SO ₂ S 6 O 2x2=4 10ve		Trigonal planar	Bent	2 Bonds 1 lone pair	Polar	1 pi 1 lone 2 sigma sp ²	120°
HCN H 1 C 4 N 5 10ve	H-C≡N:	Linear	Linear	2 Bonds 0 lone pair (only concerned w/ center)	Polar	2 pi 2 sigma sp	180
PF ₅ P 5 F 5x1=5 10ve		trigonal bi pyramid	trigonal Bi pyramid	5 Bonds 0 lone pair	NON POLAR	5 Bonds dsp ³	90/120
SCl ₄ S 6 Cl 4x1=4 10ve 4x8=32 2 lone pair		trigonal Bi pyramid	See saw	4 Bonds 1 lone pair	Polar	5 Bonds Total dsp ³	90/120
XeFCl ₃ Xe 8 F 7 Cl 3x7=21 36ve 4x8=32 2 lone pair		octahedral	Square Planar	4 Bonds 2 lone pair	δ^- F 4.0 δ^+ Cl 3.0 cancel	6 Bonds d ² sp ³	90°

4x8=32 ∴ 2 lone pair

c) Identify any molecules above that has pi bonding?

SO₂ Polar

Key

4. A compound consists of 61.70 %Cl 10.40 %C and 27.80 %S by mass. Knowing this determine the following.

a. What is the empirical formula (which is also the molecular formula) of this compound?

$$\left(\frac{61.70 \text{ g Cl}}{1}\right) \left(\frac{1 \text{ mole Cl}}{35.45 \text{ g Cl}}\right) = 1.740 / .8659 \approx 2$$

$$\left(\frac{10.40 \text{ g C}}{1}\right) \left(\frac{1 \text{ mole C}}{12.01 \text{ g C}}\right) = .8659 / .8659 \approx 1$$


$$\left(\frac{27.80 \text{ g S}}{1}\right) \left(\frac{1 \text{ mole S}}{32.07 \text{ g S}}\right) = .8688 / .8659 \approx 1$$



b. Draw the Lewis dot structure. (Note: Carbon is the central atom!)

C 4
S 6
Cl 2x7 = 14


24ve



c. This molecule has 3 sigma and 1 pi bonds.

d. What is the hybridization around the carbon? sp²

3 sigma
1 pi




Key

5. Carbon dioxide is bubbled into water.

a. Write a net ionic equation for this reaction.



b. Draw the Lewis structures of the reactants and products. Include any valid resonance structures.



No Resonance due to H's & support w/ formal charges

c. Given the following bond enthalpies, estimate the enthalpy of the reaction.

BOND	BOND ENTHALPY (kJ/mol)
H - H	436
H - O	463
O - O	146
C - O	358
C = O	799
C O	1072


$$\begin{aligned} \Delta H &= \sum H_B - \sum H_P \\ &= 2(\text{C}=\text{O}) + 2(\text{O}-\text{H}) - [2(\text{O}=\text{C}) + 2(\text{C}-\text{O}) + 2(\text{O}-\text{H})] \\ &= 2(799) + 2(463) - [2(799) + 2(358) + 2(463)] \\ &= 2524 \frac{\text{kJ}}{\text{mole}} - 2441 \frac{\text{kJ}}{\text{mole}} \end{aligned}$$

$$\Delta H = 83 \frac{\text{kJ}}{\text{mole}}$$

d. Are the C - O bonds in the reactant stronger or weaker than those in the product? Explain. Is your explanation consistent with the sign of the enthalpy change you estimated? Explain.

There are 2 x C=O Bonds to Break in CO₂ (Reactant side) Versus just one C=O in H₂CO₃. So more energy is Required to Break Bonds than produced. ∴ the Rxn should Be Endothermic & it is

e. Excess aqueous sodium hydroxide is added to the solution. Write a balanced net ionic equation for this reaction.



6. Complete the following table below.

Molecule or Ion	Lewis Dot Structure	Formal Charge on Central atom	Obeys Octet Rule Yes or No	Polar Yes or No	Resonance Yes or No
SiO_3^{2-} Si 4 $3 \times 6 = 18$ $2e = 2$ <hr/> $24ve$ $3 \times 8 = 24$		$\begin{array}{r} \text{Si} \\ 4ve \\ -4 \text{ bonds} \\ \hline 0 \end{array}$	yes	NO	yes
HCO_3^- H 1 C 4 $3 \times 6 = 18$ $1e = 1$ <hr/> $24ve$		$\begin{array}{r} \text{C} \\ 4 \\ -4 \\ \hline 0 \end{array}$	yes	yes	yes
SeO_2 Se 6 $2 \times 6 = 12$ <hr/> $18ve$		$\begin{array}{r} \text{Se} \\ 6 \\ -5 \\ +1 \end{array}$	yes	yes	yes
COS (carbon is the central atom) C 4 O 6 S 6 <hr/> $16ve$	$\begin{array}{c} \text{O} \quad \text{O} \quad \text{O} \\ \text{O} = \text{C} = \text{S} \\ \text{O} = \text{C} - \text{S} \\ \text{O} - \text{C} = \text{S} \end{array}$ <p>Yes</p>	$\begin{array}{r} \text{C} \\ 4 \\ -4 \\ \hline 0 \end{array}$	yes	$\begin{array}{r} 0 \quad 5 \\ 3.5 \quad -2.5 = 1.0 \end{array}$ Polar yes $\delta^- \text{O} = \text{S} \delta^+$	NO